

with next communication to applicant.



## U.S. DEPARTMENT OF COMMERCE ATTY DOCKET NO. CAMPBELL=2C FORM PTO-1449 PATENT AND TRADEMARK OFFICE LIST OF DOCUMENTS CITED BY APPLICANT APPLICANT: Robert CAMPBELL et al. (Use several sheets if necessary) FILING DATE: December 9, 2001 GROUT U.S. PATENT DOCUMENTS (include at least patentee, patent number, and issue date) **EXAMINER** DOCUMENT NUMBER DATE **PATENTEE CLASS SUBCLASS** FILING DATE IF INITIAL APPROPRIATE 9 CAPON et al. 435 AA 6 6 4 5/26/92 69 7 ΑB 1 5 5 0 2 7 10/13/92 SLEDZIEWSKI et al. 69 7 5 7 69.51 AC 1 1 10/22/96 RALPH et al. 12/12/94 6 AD 5 7 0 5 4 7 8 1/6/98 BOIME 11/4/94 ΑE 5 0 5 0 7/22/97 **GILLIES** 134.1 6 5 1 7/27/94 ΑF 4 4 7 8 5 1 9/5/95 BEUTLER et al. FOREIGN PATENT DOCUMENTS (include at least document number, publication date and country) TRANSLATION DOCUMENT NUMBER COUNTRY **CLASS SUBCLASS** YES DATE NO AG WO 95/31544 23NO1993 WIPO ΑH WO 9319777 14OC1993 **PCT** OTHER DOCUMENTS (include at least document number, publication date and country) AB NARAYAN, Prema et al., "Functional expression of yoked human chorionic gonadotropin in baculovirus-infected ΑI insect cells.", MOLECULAR ENDOCRINOLOGY, vol. 9, no. 12, pp. 1719-1726 (1995). JOHNSON, Gregory A. et al., "Baculovirus-insect cell production of bioactive choriogonadotropin-ΑJ immunoglobulin G heavy-chain fusion proteins in sheep.", BIOLOGY OF REPRODUCTION, vol. 52, pp. 68-73 (1995) WU, Chengbin et al., "Protein engineering of a novel constitutively active hormone-receptor complex.", JOURNAL OF ΑK BIOLOGICAL CHEMISTRY, vol. 271, no. 49, pp. 31638-31642 (1996). SMITH, Richard et al., "The active form of tumor necrosis factor is a trimer.", JOURNAL OF BIOLOGICAL ALCHEMISTRY, col 262, no. 15, pp. 6951-6954 (1987). ECK, Michael et al., "The structure of tumor necrosis factor-alpha at 2.6 A resolution.", JOURNAL OF BIOLOGICAL AM CHEMISTRY, vol. 264, no. 29, pp. 17595-17605 (1989). AN JONES, E.Y. et al., "Structure of tumor necrosis factor.", NATURE, vol. 338, pp. 225-228 (1989). ECK, Michael et al., "The structure of human lymphotoxin (tumor necrosis factor-beta) at 1.9 A resolution." JOURNAL AO OF BIOLOGICAL CHEMISTRY, vol. 267, no. 4, pp. 2119-2122 (1992). AP PIERCE, John et al., "Glycoprotein hormones structure and function.", DEPARTMENT OF BIOLOGICAL CHEMISTRY, Univ. of Cal., (1991). DATE CONSIDERED **EXAMINER EXAMPNER:** Initial if reference considered. Draw line through citation if not in conformance and not considered. Include copy of this form

ATTY DOCKET NO. U.S. DEPARTMENT OF COMMERCE FORM PTO-1449 PATENT AND TRADEMARK OFFICE CAMPBELL=2C Assigned LIST OF DOCUMENTS CITED BY APPLICANT APPLICANT: Robert CAMPBELL et al. (Use several sheets if necessary) GROUP: FILING DATE: December 9, 2001 **EXAMINER** OTHER DOCUMENTS (include at least document number, publication date and country) **INITIALS** LAPTHORN, A.J. et al., "Crystal structure of human chorionic gonadotropin.", NATURE, vol. 369, pp. 455-461 (1994). AQ AR WU, Hao et al.. "Structure of human chorionic gonadotropin at 2.6-A resolution from MAD analysis of the selenomethionyl protein.", STRUCTURE, vol. 2, no. 6, pp. 545-558 (1994). AS ENGLEMANN, Hartmut et al., "Antibodies to a soluble form of a tumor necrosis factor (TNF) receptor have TNF- like activity.", JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 265, no. 24, pp. 14497-14504 (1990). ΑT ADAM, Dieter et al., "Cross-linking of the p55 tumor necrosis factor receptor cytoplasmic domain by a dimeric ligand induces nuclear factor-Kbeta and mediates cell death.", JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 29, pp. 17482-17487 (1995). LOETSCHER, Hansruedi et al., "Recombinant 55-kda tumor necrosis factor (tnf) receptor.", JOURNAL OF ΑU BIOLOGICAL CHEMISTRY, col. 266, no. 27, pp. 18324-18329 (1991). BANNER, David W. et al., "Crystal structure of the soluble human 55kd tnf receptor-human tnfbeta complex: ΑV implications for tnf receptor activation.", CELL, vol. 73, pp. 431-445 (1993). PENNICA, Diane et al., "Biochemical characterization of the extracellular domain of the 75-kilodalton tumor necrosis AW factor receptor.", BIOCHEMISTRY, vol. 32, pp. 3131-3138 (1993). ENGELMANN, Hartmut et al., "Two tumor necrosis factor-binding proteins purified from human urine.", JOURNAL OF AXBIOLOGICAL CHEMISTRY, vol. 265, no. 3, pp. 1531-1536 (1990). VAN ZEE, Kimberly et al., "Tumor necrosis factor soluble receptors circulate during experimental and clinical ΑY inflammation and can protect against excessive tumor necrosis factor aolha in vitro and in vivo." PROC. NATL. ACAD. SCI., col. 89, pp. 4845-4849 (1992). ADERKA, Dan et al., "Stabilization of th bioactivity of tumor necrosis factor by its soluble receptors.", J. EXP. MED., AZcol. 175, pp. 323-329 (1992). MOHLER, Kendall et al., "Soluble tumor necrosis factor (tnf) receptors are effective therapeutic agents in lethal BA endotoxemia and function simultaneously as both tnf carriers and tnf antagonists.", JOURNAL OF IMMUNOLOGY, vol. 151, no. 3, pp. 1548-1561 (1993). BBBERTINI, Riccardo et al., "Urinary tnf-binding protein (tnf soluble receptor) protects mice against the lethal effect of tnf and endotoxic shock.", EUR. CYTOKINE NETW., vol. 4, no. 1, pp. 39-42 (1993). BC PIGUET, P.F. et al., "Evolution of collagen arthritis in mice is arrested by treatment with anti-tumor necrosis factor (tnf) antibody or a recombinant tnf receptor.", IMMUNOLOGY, vol. 77, pp. 510-514 (1992). WILLIAMS, Richard et al., "Successful therapy of collagen-induced arthritis with tnf receptor-IgG fusion protein and BD combination with anti-CD4.", IMMUNOLOGY, vol. 84, pop. 433-439 (1995). CAPON, Daniel et al., "Designing CD immunoadhesions for AIDS therapy." NATURE, vol. 337, pp. 525-531 (1989) BE EXAMINER DATE CONSIDERED

**EXAMPLE:** Initial if reference considered. Draw line through citation if not in conformance <u>and</u> not considered. Include copy of this form with next communication to applicant.





## FORM PTO-1449

## U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTY DOCKET NO.

CAMPBELL=2C

SERIAL NO. 09/456/86 Not Yet Assigned

LIST OF DOCUMENTS CITED BY APPLICANT

(Use several sheets if necessary)

APPLICANT: Robert CAMPBELL et al.

FILING DATE: December 9, 2001 | GROUP:

**EXAMINER** OTHER DOCUMENTS (include at least document number, publication date and country) **INITIALS** BF ASHKENZAZI, Avi et al., "Protection against endotoxic shock by a tumor necrosis factor receptor immunoadhesion.", PROC. NATL. ACAD. SCI. USA, vol. 88, pp. 10535-10539 BG SUITTERS, Amanda et al., "Differential effect of isotype on efficacy of anti-tumor necrosis factor alpha chimeric antibodies in experimental septic shock.", J. EXP. MED., vol. 179, pp. 849-856 (1994). NOLAN, Orla et al., "Bifunctional antibodies: concept, production and applications.", BIOCHIMICA ET BIOPHYSICA BH ACTA, vol. 1040, pp. 1-11 (1990). ΒI RODRIQUES, Maria L. et al., "Engineering Fab' fragments for efficient F9ab)2 formation in escherichia coli and for improved in vivo stability.", JOURNAL OF IMMUNOLOGY, vol. 151, no. 12, pp. 6954-6961 (1993). CHANG, Hsiu-Ching et al., "A general method for facilitating heterodimeric pairing between two proteins: application to BJ expression of alpha and beta t-cell receptor extracellular segments.", PROC. NATL. ACAD. SCI. USA, vol. 91, pp. 11408-11412 (1994). KIRK, Zining et al., "Solution assembly of a soluble, heteromeric, high affininty interleukin-2 receptor complex.", BK JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 27, pp. 16039-16044 (1995). BAZZIONI, F. et al., "Chimeric tumor necrosis factor receptors with constitutive signaling activity.", PROC. NATL. BLACAD. SCI. USA, vol. 92, pp. 5376-5380 (1995). BMBOLDIN, Mark et al., "Self-association of the "Death domains" of the p55 tumor necrosis factor (tnf) receptor and fas/apol prompts signaling for tnf and fas/apol effects.", JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 270, no. 1, pp. 387-391 (1995). BN VU, Thien Khai et al., "Molecular cloning of a functional thrombin receptor reveals a novel proteolytic mechanism of receptor activation.", CELL, vol. 64, pp. 1057-1068 (1991). во SONG, Ho Yeong et al., "Aggregation of the intracellular domain of the type I tumor necrosis factor receptor defined by the two-hybrid system.", J. OF BIOLOGICAL CHEMISTRY, vol. 269, no. 36, pp. 22492-22495 (1994). BP RUSSELL, Deborah et al., "Combined inhibition of interleukin-1 and tumor necrosis factor in rodent endotoxemia: improved survival and organ function.", J. INFECTIOUS DISEASES, vol. 171, pp. 1528-38 (1995). BQ RAO, Ch. et al., "Stability of human chorionic gonadotropin and its alpha subunit in human blood.", AM. J. OBSTET. GYNECOL., vol. 146, no. 1, pp. 65-68 (1983). BR DAMEWOOD, Marian et al., "Disappearance if exogenously administered human chorionic gonadotropin.", FERTILITY AND STERILITY, vol. 52, no. 3, pp. 398-400 (1989). BS CHEN, Fang et al., "The carboxy-terminal region of the glycoprotein hormone alpha-subunit: contributions to receptor binding and signaling in human chorionis gonadotropin.", MOLECULAR ENDOCRINOLOGY, vol. 6. (1992). Abstract of BIELINSKA, M. et al., "Site-directed mutagenesis identifies two receptor binding domains in the human BT chorionic gonadotropin alpha subunit.", MEMBRANE RECEPTORS, no. 1844. BU FURUHASHI, Madoka et al., "Fusing the carboxy-terminal peptide of the chronic gonadotropin (cg) beta-subunit to the common alpha-subunit: retention of o-linked glycosylation and enhanced in vivo bioactivity of chimeric human cg.", MOLECULAR EMDOCRINOLOGY, 9:(1) pp. 63 (1995).

EXAMINER

J. Specto

DATE CONSIDERED

**EXAMINEF:** Initial if reference considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.





## U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

ATTY DOCKET NO.

CAMPBELL=2C

LIST OF DOCUMENTS CITED BY APPLICANT (Use several sheets if necessary)

APPLICANT: Robert CAMPBELL et al.

FILING DATE: December 9, 2001

GROUP:

		individual straight s
EXAMINER INITIALS		OTHER DOCUMENTS (include at least document number, publication date and country)
$\mathcal{J}$	BV	SUGAHRA, Tadashi et al., "Biosynthesis of a biologically active single peptide chain containing the human common alpha and chorionic gonadotropin beta subunits in tandem.", PROC. NATL. ACAD. SCI. USA, vol. 92, pp. 2041-2045 (1995).
	BW	URLAUB, Gail et al., "Isolation of chinese hamster cell mutants deficient in dihydrofolate reductase activity.", PROC. NATL. ACAD. SCI. USA, vol. 77, no. 7, pp. 4216-4220 (1980).
	BX	NOPHAR, Yaron et al., "Soluble forms of tumor necrosis factor receptors (tnf-rs).", THE EMBO JOURNAL, vol. 9, no 10, pp. 3269-3278 (1990).
	BY	FIDDES, John et al., "Isolation, cloning and sequence analysis of the cDNA for the alpha-subunit of human chorionic gonadotropin.", NATURE, vol. 281, pp. 351-356 (1979).
	BZ	FIDDES, John et al., "The cDNA for the beta-subunit of human chorionic gonadotropin suggests evolution of a gene by readthrough into the 3'-untranslated region.", NATURE, vol. 286, (1980).
	CA	CAMPBELL, Robert et al., "Conversion of human choriogonadotropin into a follitropin by protein engineering." PROC NATL. ACAD. SCI. USA, vol. 88, pp. 760-764 (1991).
	СВ	COLE, Edward et al., "Recombinant human thyroid stimulating hormone: development of a biotechnology product for detection of metastatic lesions of thyroid cancer.", BIOTECHNOLOGY, vol. 11, pp. 1014-1024 (1993).
	СС	GLUZMAN, Yakov. "SV40-transformed simian cells support the replication of early SV40 mutants.", CELL, vol. 23, p 175-182 (1981).
	CD	CHU, Gilbert., "Electroporation for the efficient transfection of mammalian cells with DNA.", NUCLEIC ACIDS RESEARCH, vol. 15, no. 3 (1987).
V	CE	YEN, Janie et al., "A rapid in vitro cytotoxicity assay for the detection of tumor necrosis factor on human BT-20 cells.", JOURNAL OF IMMUNOTHERAPY, vol. 10, pp. 174-181 (1991).
EXAMINER	<u> </u>	DATE CONSIDERED

FORM PTO-1449

**EXAMINER:** Initial if reference considered. Draw line through citation if not in conformance and not considered. Include copy of this form with next communication to applicant.